

How to Flourish in Social Work



Preserving personal longevity and happiness, relationships, and your career

COMMON AILMENTS

"A feeling of depletion leads to dysfunction... a [Social Work] practitioner becomes increasingly 'inoperative'."



BURNOUT



COMPASSION FATIGUE



SECONDARY TRAUMATIC STRESS

STEPS TO SELF-CARE

Self-care refers to selected actions that restore balance in our personal and professional lives.

Not just an add-on activity, self-care is also a state of mind through development of self-awareness, self-regulation, and self-efficacy.



EXERCISE

Light **3 DAYS A WEEK** exercise improves happiness by **10-20%** increases work productivity by **15%**



MEDITATE

50% reduction in overall psychiatric symptoms
70% decrease in anxiety
44% reduction in common medical symptoms



READ

A new trend in treatment of mental illness; boosts creativity and activates sensory areas of the brain.



GREENSPACE

Having **over 30%** of green space in your surroundings is recommended for **healthy cortisol levels**.



LAUGH

Laughter strengthens the immune system, boosts energy and diminishes pain.

Children laugh over **300 times per day**
The typical adult chuckles **15 times per day**



TIME OFF

30% of employees use their vacation time, which leads to better quality sleep, decreased stress and improved mood.



EAT WELL

Omega-3 fatty acids improve learning and memory and fight mental disorders. Carbohydrates aid in the release of endorphins.



SLEEP

The CDC currently classifies insufficient sleep as a public health epidemic. Sleep restores cognitive functions.



For a self-care starter kit, please visit
<http://www.socialwork.buffalo.edu/students/self-care>

SCHOOL OF SOCIAL WORK
University at Buffalo The State University of New York

Adams, R. E., Faye, C. R., & Brannigan, J. A. (2008). The connection between social work and social workers following urban disaster. *Disaster Prevention and Management: An International Journal, 19*(2), 218-230. doi:10.1108/09697530810873103

Baker, S. S. (2007). Prevalence of secondary traumatic stress among social workers. *Social Work, 52*, 43-50.

Center for Workplace Studies. (n.d.). Retrieved from <http://www.cwps.buffalo.edu/>

Carroll, T. A. (1998). Secondary traumatic stress among child protective service workers. *Social Work, 43*, 10-18. doi:10.1177/002615419804300108

National Alliance on Mental Illness. (2014). Retrieved from <http://www.nami.org/>

Smolkin, Sara Kay. "What I Wish I Had Known About Self-Care in Our Social Work Professions." *The New Social Worker*, Fall 2012. Retrieved from <http://www.nswa.org.uk/>

Soko, S. (2009). Reflections on the conceptualization of compassion fatigue. *Omega: The Official Journal of Issues in Learning, 47*(1), doi:10.1016/j.omega.2008.11.001

United States Census Bureau. (2014). Retrieved from <http://www.census.gov/>